

Nodal solutions of a NLS equation concentrating on lower dimensional spheres

Giovany M. Figueiredo*
Faculdade de Matemática
Universidade Federal do Pará
66075-110, Belém - Pa, Brazil.
e-mail: giovany@ufpa.br

and

Marcos T. O. Pimenta^{†‡}
Departamento de Matemática e Computação
Faculdade de Ciências e Tecnologia
Universidade Estadual Paulista - Unesp
19060-900, Presidente Prudente - SP, Brazil.
e-mail: pimenta@fct.unesp.br

Abstract

In this work we deal with the following nonlinear Schrödinger equation

$$\begin{cases} -\epsilon^2 \Delta u + V(x)u = f(u) & \text{in } \mathbb{R}^N \\ u \in H^1(\mathbb{R}^N), \end{cases}$$

where $N \geq 3$, f is a subcritical power-type nonlinearity and V is a positive potential satisfying a local condition. We prove the existence

*Supported by PROCAD/CASADINHO 552101/2011-7, CNPq/PQ 301242/2011-9 and CNPq/CSF 200237/2012-8

[†]Corresponding author

[‡]Supported by FAPESP 2014/16136-1 and CNPq 442520/2014-0

and concentration of nodal solutions which concentrate around a k -dimensional sphere of $\mathbb{R}^N$, where $1 \leq k \leq N - 1$, as $\epsilon \rightarrow 0$. The radius of such sphere is related with the local minimum of a function which takes into account the potential V . Variational methods are used together with the penalization technique in order to overcome the lack of compactness.

2010 Mathematics Subject Classification: 35J60, 35J10, 35J20.

Keywords: variational methods, nodal solutions, concentration on manifolds.

1 Introduction

In the last decades, motivated by the great interest that this problem catch in quantum mechanics, so many researchers have dedicated their efforts on the study of the Nonlinear Schrödinger equation

$$i\hbar \frac{\partial \psi}{\partial t} = -\frac{\hbar^2}{2m} \Delta \psi + W(x)\psi - |\psi|^{p-1}\psi \quad (t, x) \in \mathbb{R} \times \mathbb{R}^N.$$

Of particular interest are the so called *standing wave solutions* which consists in solutions with a particle-like behavior. It is obtained by the *Ansatz* $\psi(t, x) = e^{-iEt/\hbar} u(x)$ which associate the NLS equation to its stationary version

$$-\epsilon^2 \Delta u + V(x)u = |u|^{p-1}u \quad \text{in } \mathbb{R}^N \quad (1.1)$$

where $\epsilon^2 = \hbar^2/2m$ and $V(x) = W(x) - E$. As far as (1.1) is concerned, the behavior of the solutions when $\epsilon \rightarrow 0$ has a great physical interest since it describes the transition from quantum to the classical mechanics, being called *semiclassical states*. On this specific subject, many authors have worked on *spike-layered solutions* which are nontrivial ground-state points of the associated energy functional and tend to concentrate around one or more critical points of the potential V . We could cite some quite influent works on this subject, as the pioneering work of Floer and Weinstein [13], which have inspired the works of Rabinowitz [18], Wang [21], Del Pino and Felmer [12], which have influenced so many other works by their own in the last three decades.

In the last ten years, solutions which concentrate on higher dimensional sets has been received more and more attention. The first work which seems to show this kind of result is [16] in which the authors study a NLS equation

on a bounded domain with Neumann boundary condition and prove the existence of a sequence of solutions which concentrate on some component of the boundary. One of the first works dealing with solutions concentrating around a sphere is [4] in which Ambrosetti, Malchiodi and Ni give necessary and sufficient conditions under which (1.1) exhibit solutions concentrating around a sphere. The radius of such a sphere is given by a minimum point of a function $\mathcal{M}$, which takes into account the value of the radial potential $V(|x|)$. The role played by $\mathcal{M}$ is in order to balance the potential energy (coming from V) and the volume energy which arise from the other terms of the energy functional (see the introduction of [4] for more details). In fact, sphere concentrating solutions show a rather different behavior when compared with spike-layered ones. To be more specific, in [4], the authors prove the existence of sphere concentrating solutions to (1.1) even for critical or supercritical exponent p . This is in a strike contrast with the fact that, as showed in [10], no spike-layered solution exists to (1.1) for $p = 2^* - 1$. Other significant difference is that the energy of the sphere concentrating solutions tend to zero, in contrast with those of spike-layered solutions, whose energy converge to the mountain-pass level of the energy functional. In these and so many other works ([5, 19, 11, 15] for example), Lyapunov-Schmidt reduction methods have been used in order to construct the sphere-concentrating solutions for Schrödinger equations, Schrödinger-Poisson systems other related problems.

More recently, in [7] Bonheure *et al* proved the existence of solutions concentrating around a k -dimensional sphere of $\mathbb{R}^N$, for all $k \in \{1, \dots, N-1\}$ to the following equation

$$-\epsilon^2 \Delta u + V(x)u = K(x)f(u) \quad \text{in } \mathbb{R}^N, \quad (1.2)$$

where the potentials V and K satisfy rather generic conditions, allowing V even to vanish on the infinity. To do so they use a modification of the penalization technique, originally presented in [12], in such a way that compactness is recovered to the modified energy functional. Because of the generality of conditions under which V and K are subjected, in order to prove that the solutions of the modified problem are solutions of the original one, they made a thorough analysis with some barrier functions which bounds the solutions from above. In [8] the authors employ a similar argument in order to show the existence of solutions concentrating on circumferences of $\mathbb{R}^3$, to a Schrödinger-Poisson system.

In the *spike-layered* solutions setting, the existence of sign-changing (or nodal) solutions was investigated by some authors. In [1] and [2], Alves and

Soares study problem (1.2), with K to be a constant, and prove the existence of nodal solutions which concentrate at minima of the potential V . In the first work they consider f as a subcritical power-type nonlinearity and in the second, as presenting a critical exponential growth at infinity. In both they employ the penalization technique together with a careful analysis of the profile of the solutions. In [20], Y. Sato have proposed a different kind of penalization in order to show the existence of multi-peak nodal solutions to a Schrödinger equation with a vanishing potential.

A question that naturally arises is whether there exist a sequence of nodal solutions to the NLS equation which concentrate around a k -dimensional sphere. In this work we give a positive answer to this question. More specifically, we study the existence and concentration of nodal solutions to the following nonlinear Schrödinger equation

$$\begin{cases} -\epsilon^2 \Delta u + V(x)u = f(u) & \text{in } \mathbb{R}^N \\ u \in H^1(\mathbb{R}^N), \end{cases} \quad (1.3)$$

where $N \geq 3$, exhibiting a cylindrical symmetry which implies in this sort of concentration. The nonlinearity f is assumed to be a $C^1(\mathbb{R})$ odd function satisfying

(f_1) There exists $\nu > 1$ such that $f(|s|) = o(|s|^\nu)$ as $s \rightarrow 0$;

(f_2) There exist $c_1, c_2 > 0$ such that $|f'(s)| \leq c_1 + c_2|s|^{p-1}$ where $1 < p < \frac{N-k+2}{(N-k-2)}$ and k is like in Section 1.1;

(f_3) There exists $\theta > 2$ such that

$$0 < \theta F(s) \leq f(s)s, \quad \text{for } s \neq 0,$$

$$\text{where } F(s) = \int_0^s f(t)dt;$$

(f_4) $s \mapsto f(s)/s$ is increasing in $s > 0$ and decreasing for $s < 0$.

The potential V will be assumed to satisfy a symmetry condition which we explain in the next section.

1.1 Statement of the main result

Let $1 \leq k \leq N - 1$ be an integer which determine the dimension of the sphere in which the solutions obtained are going to concentrate. Consider $\mathcal{H}$ a $(N - k - 1)$ -dimensional linear subspace of $\mathbb{R}^N$ and note that $\mathcal{H}^\perp$ is a $(k + 1)$ -dimensional subspace. All along the paper we use the notation for $x \in \mathbb{R}^N$ as $x = (x', x'')$ in which $x' \in \mathcal{H}$, $x'' \in \mathcal{H}^\perp$ are such that $x = x' + x''$.

From now on, if $h : \mathbb{R}^N \rightarrow \mathbb{R}$ is a function, by saying that $h(x', x'') = h(x', |x''|)$ (which rigorously does not make sense), we mean that $h(x', y) = h(x', z)$ for all $y, z \in \mathcal{H}^\perp$ such that $|y| = |z|$.

The condition in V which is considered is the following:

(V₁) There exists $V_0 > 0$ such that $V_0 \leq V(x)$ and, for all $x \in \mathbb{R}^N$, $V(x) = V(x', x'') = V(x', |x''|)$.

Unlike spike layered solutions, whose concentration occurs around minimum points of V , the solutions we are going to study concentrate around minimum points of an auxiliary potential. To see how we define it, let us consider the limit problem

$$-\Delta u + au = f(u) \quad \text{in } \mathbb{R}^{N-k}. \quad (1.4)$$

It is well known (see [18] for instance) that there exists a ground state solution $w \in H^1(\mathbb{R}^{N-k})$ of (1.4) which minimizes the energy functional

$$I_a(u) = \frac{1}{2} \int_{\mathbb{R}^{N-k}} (|\nabla u|^2 + au^2) dx - \int_{\mathbb{R}^{N-k}} F(u) dx,$$

in the corresponding Nehari manifold given by

$$\mathcal{N}_a = \{u \in H^1(\mathbb{R}^{N-k}) \setminus \{0\}; I'_a(u)u = 0\}.$$

We define the *ground-energy function* $\mathcal{E} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, by

$$\mathcal{E}(a) = \inf_{\mathcal{N}_a} I_a.$$

Finally, we define the *auxiliary potential* $\mathcal{M} : \mathbb{R}^N \rightarrow (0, +\infty]$ by

$$\mathcal{M}(x) = |x''|^k \mathcal{E}(V(x))$$

where $x = (x', x'')$, $x' \in \mathcal{H}$ and $x'' \in \mathcal{H}^\perp$.

On the auxiliary potential $\mathcal{M}$ we impose the following condition

($\mathcal{M}_1$) There exists an open bounded set $\Omega \subset \mathbb{R}^N$ such that, if $(x', x'') \in \Omega$ then $(x', y'') \in \Omega$ for all $y'' \in \mathcal{H}^\perp$, $|x''| = |y''|$. Moreover

$$0 < \mathcal{M}_0 := \inf_{x \in \Omega} \mathcal{M}(x) < \inf_{x \in \partial\Omega} \mathcal{M}(x).$$

Now we can finally state our main result.

Theorem 1.1. *Let f satisfying $(f_1) - (f_4)$ and V such that (V_1) and $(\mathcal{M}_1)$ hold. Then for each sequence $\epsilon_n \rightarrow 0$, there exists a subsequence still denoted by (ϵ_n) such that (1.3) (with $\epsilon = \epsilon_n$) has a nodal bound state u_n such that, $u(x', x'') = u(x', |x''|)$ and, if $\epsilon_n P_n^1$ and $\epsilon_n P_n^2$ are respectively a minimum and a maximum point of u_n , then $\epsilon_n P_n^i \in \Omega$, $i = 1, 2$ for n sufficiently large,*

$$\epsilon_n P_n^i \rightarrow x_0, \quad \text{as } n \rightarrow \infty \quad (1.5)$$

where $\mathcal{M}(x_0) = \mathcal{M}_0$ and

$$|u_n(x)| \leq C \left(e^{-\frac{\beta}{\epsilon_n} d_k(x, \epsilon_n P_n^1)} + e^{-\frac{\beta}{\epsilon_n} d_k(x, \epsilon_n P_n^2)} \right) \quad x \in \mathbb{R}^N,$$

where $C, \beta > 0$ and d_k is the distance defined in (4.11).

The arguments in proving the existence of solutions were strongly influenced by the works of Alves and Souto [3], in which they prove the existence of nodal solutions to a Schrödinger-Poisson system. In the concentration, we follow closely the arguments in [1] and [7, 8].

After our work has been finished we found the very recent paper [14] in which the author uses a similar argumentation in order to prove the existence of a sequence of nodal multi-peak solutions which concentrate around the minimum points of a modified potential, associated to a vanishing potential. The existence arguments in both works rely on a minimization of the penalized energy functional on the nodal Nehari set and the concentration arguments follow the same general lines. Nevertheless, it is worth pointing out that in our work, since we get sphere concentrating solutions, several technical difficulties arise. Moreover, in our work proving that the solution of the modified problem is in fact a solution of the original one involves different comparison functions, since our penalization is slightly different.

In Section 2 we present the penalization scheme and the variational framework. In Section 3 we prove the existence of the nodal solutions of the modified problem. In Section 4 we exhibit the concentration arguments

in order to prove that the solutions of the modified problem concentrates around a k -dimensional sphere and in the last section we complete the prove of the Theorem 1.1 by showing that the solutions of the modified problem satisfy the original one.

2 The penalized nonlinearity and the variational framework

The penalization we are going to apply is a variation of the classical method of Del Pino and Felmer in [12], developed by Y. Sato in [20] in order to allow its use in finding nodal solutions. Fixing $2 < \tau < \theta$, let $r_\epsilon > 0$ such that

$$\frac{f(r_\epsilon)}{r_\epsilon} = \epsilon^\tau \quad \text{and} \quad \frac{f(-r_\epsilon)}{-r_\epsilon} = \epsilon^\tau.$$

Since $r_\epsilon \rightarrow 0$ as $\epsilon \rightarrow 0$, (f_1) implies that

$$\epsilon^\tau = \frac{f(|r_\epsilon|)}{|r_\epsilon|} \leq |r_\epsilon|^{\nu-1}.$$

Thus $\epsilon^{\frac{\tau}{\nu-1}} \leq |r_\epsilon|$ and we can choose an odd function $\tilde{f}_\epsilon \in C^1(\mathbb{R})$ satisfying

$$\tilde{f}_\epsilon(s) = \begin{cases} f(s) & \text{if } |s| \leq \frac{1}{2}\epsilon^{\frac{\tau}{\nu-1}}, \\ \epsilon^\tau s & \text{if } |s| \geq \epsilon^{\frac{\tau}{\nu-1}}, \end{cases}$$

$$|\tilde{f}_\epsilon(s)| \leq \epsilon^\tau |s| \quad \text{for all } s \in \mathbb{R}, \quad (2.1)$$

$$0 \leq \tilde{f}'_\epsilon(s) \leq 2\epsilon^\tau \quad \text{for all } s \in \mathbb{R} \quad (2.2)$$

and

$$s \mapsto \tilde{f}_\epsilon(s)/s \text{ is increasing for } s > 0 \text{ and decreasing for } s < 0. \quad (2.3)$$

Let us define $g_\epsilon(x, s) := \chi_\Omega(x)f(s) + (1 - \chi_\Omega(x))\tilde{f}_\epsilon(s)$, where χ_Ω is the characteristic function of Ω . Note that by $(f_1) - (f_4)$, g is a Charathéodory function, such that $g_\epsilon(x', x'', s) = g_\epsilon(x', |x''|, s)$ satisfying

(g_1) $g_\epsilon(x, s) = o(|s|^\nu)$, as $s \rightarrow 0$, uniformly in compact sets of $\mathbb{R}^N$.

(g_2) There exist $c_1, c_2 > 0$ such that $|g_\epsilon(x, s)| \leq c_1|s| + c_2|s|^p$ where $1 < p < \frac{N+2}{N-2}$;

(g_3) There exists $\theta > 2$ such that:

$$i) \quad 0 < \theta G_\epsilon(x, s) \leq g_\epsilon(x, s)s, \text{ for } x \in \Omega \text{ and } s \neq 0,$$

$$ii) \quad 0 < 2G_\epsilon(x, s) \leq g_\epsilon(x, s)s, \text{ for } x \in \mathbb{R}^N \setminus \Omega \text{ and } s \neq 0,$$

$$\text{where } G_\epsilon(x, s) = \int_0^s g_\epsilon(x, t)dt.$$

(g_4) $s \mapsto \frac{g_\epsilon(x, s)}{s}$ is a nondecreasing function for $s > 0$ and nonincreasing for $s < 0$, for all $x \in \mathbb{R}^N$.

In a first moment, the concern will be with the penalized problem

$$-\epsilon^2 \Delta u + V(x)u = g_\epsilon(x, u) \quad \text{in } \mathbb{R}^N. \quad (2.4)$$

Taking $v_\epsilon(x) = u_\epsilon(\epsilon x)$, we relate each solution u_ϵ of (2.4) with a solution v_ϵ of

$$-\Delta v + V(\epsilon x)v = g_\epsilon(\epsilon x, u) \quad \text{in } \mathbb{R}^N. \quad (2.5)$$

In order to obtain solutions of (2.5) with a partial symmetry, let us consider the following subspace of $H^1(\mathbb{R}^N)$,

$$\tilde{H} := \left\{ v \in H^1(\mathbb{R}^N); \int (|\nabla v|^2 + V(\epsilon x)v^2) < +\infty \text{ and } v(x', x'') = v(x', |x''|) \right\}$$

which is a Hilbert space when endowed with the inner product

$$\langle u, v \rangle_\epsilon = \left(\int (\nabla u \nabla v + V(\epsilon x)uv) \right),$$

which gives rise to the following norm

$$\|v\|_\epsilon = \left(\int (|\nabla v|^2 + V(\epsilon x)v^2) \right)^{\frac{1}{2}}.$$

Since the approach is variational, let us consider the energy functional $I_\epsilon : \tilde{H} \rightarrow \mathbb{R}$, whose Euler-Lagrange equation is (2.5), given by

$$I_\epsilon(v) = \frac{1}{2} \int (|\nabla v|^2 + V(\epsilon x)v^2) - \int G_\epsilon(\epsilon x, v).$$

By standard arguments, one can prove that $I_\epsilon \in C^2(\tilde{H}, \mathbb{R})$.

Remark 2.1. In this section and through the rest of the paper, we omit the dx in all the integrals and, when the domain over which the integral is calculated is $\mathbb{R}^N$ we write $\int$ rather than $\int_{\mathbb{R}^N}$.

3 Existence results

Let us consider the Nehari manifold associated to (2.5), which is well defined by (g_4) and given by

$$\mathcal{N}_\epsilon = \{v \in \tilde{H} \setminus \{0\}; I'_\epsilon(v)v = 0\}.$$

Since we are looking for nodal solutions, let us consider the so called nodal Nehari set

$$\mathcal{N}_\epsilon^\pm = \{v \in \tilde{H}; v^\pm \neq 0 \text{ and } I'_\epsilon(v)v^\pm = 0\}.$$

Although $\mathcal{N}_\epsilon^\pm$ is not a manifold since $u \mapsto u^\pm$ in $H^1(\mathbb{R}^N)$ lacks differentiability, it is a set which contain all nodal solutions of (2.5).

Next result try to infer informations of I_ϵ with respect to $\mathcal{N}_\epsilon^\pm$ in the same way that one is used to do with $\mathcal{N}_\epsilon$.

Lemma 3.1. *Let $v \in \tilde{H}$ such that $v^\pm \neq 0$. Then there exist $t, s > 0$ such that $tv^+ + sv^- \in \mathcal{N}_\epsilon^\pm$.*

Proof. First of all let us prove that, for all $v \in \tilde{H} \setminus \{0\}$, there exists $t > 0$ such that $tv \in \mathcal{N}_\epsilon$. Indeed, if $\Gamma = (\text{supp } v \cap \Omega_\epsilon) \cup \{x \in \mathbb{R}^N; |v(x)| \leq \frac{1}{2}\epsilon^{\frac{\tau}{\tau-1}}\}$, note that $|\Gamma| > 0$ (since $v \in H^1(\mathbb{R}^N)$) and then

$$\begin{aligned} I_\epsilon(tv) &= \frac{t^2}{2} \|v\|_\epsilon^2 - \int_\Gamma F(tv) - \int_{\mathbb{R}^n \setminus \Gamma} G(\epsilon x, v) \\ &\leq \frac{t^2}{2} \|v\|_\epsilon^2 - t^\theta \int_\Gamma |v|^\theta - \frac{t^\tau}{2} \int_{\mathbb{R}^n \setminus \Gamma} |v| \\ &\rightarrow -\infty, \end{aligned}$$

as $t \rightarrow \infty$. Then, if $v \in \tilde{H}$ is such that $v^\pm \neq 0$, there exist $t, s > 0$ such that

$$I'_\epsilon(tv^+)tv^+ = 0 \quad \text{and} \quad I'_\epsilon(sv^-)sv^- = 0.$$

Then, it is clear that

$$I'_\epsilon(tv^+ + sv^-)(tv^+ + sv^-) = I'_\epsilon(tv^+)tv^+ + I'_\epsilon(sv^-)sv^- = 0.$$

□

For a fixed $v \in \tilde{H}$, let us consider $\psi_v : [0, +\infty) \times [0, +\infty) \rightarrow \mathbb{R}$ given by

$$\psi_v(t, s) = I_\epsilon(tv^+ + sv^-),$$

and note that by the smoothness of g , $\psi_v \in C^2(\mathbb{R}^2, \mathbb{R})$.

Lemma 3.2. *Let $v \in \mathcal{N}_\epsilon^\pm$, then $(t, s) = (1, 1)$ is a strict global maximum point of ψ_v .*

Proof. First of all let us note that if $v \in \mathcal{N}_\epsilon^\pm$, by (g_3)

$$\lim_{|(t,s)| \rightarrow \infty} \psi_v(t, s) = -\infty.$$

Then there exists $R > 0$ such that

$$\psi_v(t, s) < 0, \quad \text{if } |(t, s)| \geq R. \quad (3.1)$$

Since

$$\nabla \psi_v(t, s) = (I'_\epsilon(tv^+)v^+, I'_\epsilon(sv^-)v^-),$$

standard calculations about the behavior of $t \mapsto I_\epsilon(tv^+)$ and $s \mapsto I_\epsilon(sv^-)$ and the fact that $v \in \mathcal{N}_\epsilon^\pm$, implies that ψ_v has just one critical point given by $(t, s) = (1, 1)$.

As we prove in Lemma 3.5 (which is totally independent of this one), $\psi_v(1, 1) = I_\epsilon(v) \geq \rho > 0$. By (3.1) in order to get the result its enough to prove that $(1, 1)$ is a local maximum point of ψ_v . Note that

$$D^2\psi_v(t, s) = \begin{pmatrix} I''_\epsilon(tv^+)(v^+, v^+) & 0 \\ 0 & I''_\epsilon(sv^-)(v^-, v^-) \end{pmatrix}$$

and then

$$\begin{aligned} \det(D^2\psi_v(1, 1)) &= I''_\epsilon(v^+)(v^+, v^+) \cdot I''_\epsilon(v^-)(v^-, v^-) \\ &= \left(\int \left(g_\epsilon(\epsilon x, v^+)v^+ - g'_\epsilon(\epsilon x, v^+)v^{+2} \right) \right) \\ &\quad \cdot \left(\int \left(g_\epsilon(\epsilon x, v^-)v^- - g'_\epsilon(\epsilon x, v^-)v^{-2} \right) \right). \end{aligned}$$

By definition of g_ϵ and (g_4) , the last integral is greater or equal to

$$\begin{aligned} &\left(\int_{(supp(v^+) \cap \Omega_\epsilon) \cup \{|v^+| \leq \frac{1}{2}\epsilon^{\frac{\tau}{\nu-1}}\}} \left(f(v^+)v^+ - f'(v^+)v^{+2} \right) \right) \\ &\cdot \left(\int_{supp(v^-) \cap (\Omega_\epsilon \cup \{|v^-| < \frac{1}{2}\epsilon^{\frac{\tau}{\nu-1}}\})} \left(f(v^-)v^- - f'(x, v^-)v^{-2} \right) \right) \\ &> 0. \end{aligned}$$

In the last inequality we have used that by (f_4) ,

$$f(s)s - f'(s)s^2 < 0, \quad \text{for all } s \neq 0$$

and as $v^\pm \in \tilde{H}$, $|supp(v^+) \cap (\Omega_\epsilon \cup \{|v^+| < a\})| > 0$ and $|supp(v^-) \cap (\Omega_\epsilon \cup \{|v^-| < a\})| > 0$, where $\Omega_\epsilon := \epsilon^{-1}\Omega$.

Since $D^2\psi_v(1,1)$ is a positive definite form, we have just to verify that $\frac{\partial^2\psi_v}{\partial t^2} = I''_\epsilon(tv^+)v^{+2} < 0$. But this follows since 1 is a maximum point of $t \mapsto I_\epsilon(tv^+)$. $\square$

Still as a consequence of the arguments employed in the construction of the Nehari manifold as in [18], it follows the following result.

Lemma 3.3. *Let $v \in \tilde{H}$ such that $v^\pm \neq 0$ and*

$$I'_\epsilon(v)v^\pm \leq 0,$$

then there exists $t, s \in (0, 1]$ such that

$$tv^+ + sv^- \in \mathcal{N}_\epsilon^\pm.$$

Proof. In fact, let $t \in \mathbb{R}$ such that $I'_\epsilon(tv^+)tv^+ = 0$. Suppose by contradiction that $t > 1$, then

$$\begin{aligned} \|v^+\|^2 &= \int \frac{g_\epsilon(\epsilon x, tv^+)v^+}{t} = \int_{v^+ > 0} \frac{g_\epsilon(\epsilon x, tv^+)v^{+2}}{tv^+} \\ &= \int_{supp(v^+) \cap (\Omega_\epsilon \cup \{|tv^+| < \frac{1}{2}\epsilon^{\frac{\tau}{\nu-1}}\})} \frac{f(tv^+)tv^{+2}}{v^+} + \\ &\quad \int_{supp(v^+) \cap (\Omega_\epsilon \cup \{|tv^+| \geq \frac{1}{2}\epsilon^{\frac{\tau}{\nu-1}}\})} \frac{g_\epsilon(\epsilon x, tv^+)v^{+2}}{tv^+} \\ &> \int_{supp(v^+) \cap (\Omega_\epsilon \cup \{|tv^+| < \frac{1}{2}\epsilon^{\frac{\tau}{\nu-1}}\})} f(v^+)v^+ \\ &\quad + \int_{supp(v^+) \cap (\Omega_\epsilon \cup \{|v^+| \geq \frac{1}{2}\epsilon^{\frac{\tau}{\nu-1}}\})} \frac{g_\epsilon(\epsilon x, v^+)v^{+2}}{v^+} \\ &= \int g_\epsilon(\epsilon x, v^+)v^+, \end{aligned}$$

which implies that $I'_\epsilon(v)v^+ > 0$, contradicting the hypothesis. The same argument applies to v^- and s . $\square$

Let us define

$$d_\epsilon := \inf_{\mathcal{N}_\epsilon^\pm} I_\epsilon,$$

and note that if there exists a solution of (2.5) in the energy level d_ϵ , then it is the solution with least energy among all nodal ones.

Now we are going to state and prove the main result of this section.

Theorem 3.4. *For sufficiently small $\epsilon > 0$, there exists a nodal solution of (2.5), $v_\epsilon \in \tilde{H}$ such that $I_\epsilon(v_\epsilon) = d_\epsilon$.*

Before proceed with the proof of Theorem 3.4 let us state some technical result about $\mathcal{N}_\epsilon^\pm$.

Lemma 3.5. *It holds that*

i) *There exists $\rho > 0$ such that $\|v\|_\epsilon \geq \rho$ for all $v \in \mathcal{N}_\epsilon$.*

ii) *There exists a constant $C > 0$ such that, for all $v \in \mathcal{N}_\epsilon$, $I_\epsilon(v) \geq C\|v\|_\epsilon^2$.*

Proof. The proof of i) follows by standard arguments. Let us prove just ii), which in fact also follows by very known arguments. Note that, if $v \in \mathcal{N}_\epsilon$, by (g_3)

$$\begin{aligned} I_\epsilon(v) = I_\epsilon(v) - \frac{1}{\theta} I'_\epsilon(v)v &= \left(\frac{1}{2} - \frac{1}{\theta} \right) \|v\|_\epsilon^2 + \frac{1}{\theta} \int_{\mathbb{R}^N} (g(x, v)v - \theta G(x, v)) dx \\ &\geq \left(\frac{1}{2} - \frac{1}{\theta} \right) \|v\|_\epsilon^2 + \frac{1}{\theta} \int_{\Omega_\epsilon^c} (g(\epsilon x, v_n)v - \theta G(\epsilon x, v)) dx \\ &\geq \left(\frac{1}{2} - \frac{1}{\theta} \right) \|v\|_\epsilon^2 + \frac{(2-\theta)}{\theta} \int_{\Omega_\epsilon^c} G(x, u_n) dx \\ &\geq \left(\frac{1}{2} - \frac{1}{\theta} \right) \|v\|_\epsilon^2 + \frac{(2-\theta)}{2\theta} \int_{\Omega_\epsilon^c} \epsilon^\tau v^2 dx \\ &= \left(\frac{\theta-2}{2\theta} \right) \int_{\mathbb{R}^N} (|\nabla v|^2 + (V(\epsilon x) - \epsilon^\tau)v^2) dx \\ &= \left(\frac{\theta-2}{2\theta} \right) \int_{\mathbb{R}^N} (|\nabla v|^2 + (V_0 - \epsilon^\tau)v^2) dx \\ &\geq C \|v\|_\epsilon^2, \end{aligned}$$

where $C > 0$ for $\epsilon > 0$ sufficiently small. □

Proof of Theorem 3.4. The proof will be carried out into two steps. In the first one we prove that d_ϵ is attained by a function $u_\epsilon \in \tilde{H}$.

Let (w_n) be a minimizing sequence for I_ϵ in $\mathcal{N}_\epsilon^\pm$, i.e., a sequence $(w_n) \subset \mathcal{N}_\epsilon^\pm$ such that

$$\lim_{n \rightarrow \infty} I_\epsilon(w_n) = d_\epsilon. \quad (3.2)$$

Note that by (3.2) and Lemma 3.5, (w_n) is a bounded sequence in $\tilde{H}$. Then there exists w_ϵ such that $w_n \rightharpoonup w_\epsilon$ in $\tilde{H}$ up to a subsequence. In the same way as in Lemma 2.3 in [9], it is possible to show that $v \mapsto v^\pm$ is a continuous function of $\tilde{H}$ into itself, from which follows that $w_n^\pm \rightharpoonup w_\epsilon^\pm$ in $\tilde{H}$. As a consequence, up to a subsequence

$$w_n^\pm \rightarrow w_\epsilon^\pm, \quad \text{a.e. in } \mathbb{R}^N. \quad (3.3)$$

and

$$w_n^\pm \rightarrow w_\epsilon^\pm, \quad \text{in } L^r(\Omega_\epsilon), \text{ for } 1 \leq r < \frac{2(N-k)}{N-k-2} \quad (3.4)$$

where $w_\epsilon^\pm \neq 0$ by the same arguments in Lemma 2.8 in [20].

Since $w_\epsilon^\pm \neq 0$, let $t_\epsilon, s_\epsilon > 0$ be such that $t_\epsilon w_\epsilon^+ + s_\epsilon w_\epsilon^- \in \mathcal{N}_\epsilon^\pm$. By weak lower-semicontinuity of $\|\cdot\|$ and by Sobolev embeddings it follows that

$$\|t_\epsilon w_\epsilon^+ + s_\epsilon w_\epsilon^-\|_\epsilon \leq \liminf_{n \rightarrow \infty} \|t_\epsilon w_n^+ + s_\epsilon w_n^-\|_\epsilon \quad (3.5)$$

and

$$\int_{\Omega_\epsilon} F(t_\epsilon w_\epsilon^+ + s_\epsilon w_\epsilon^-) = \liminf_{n \rightarrow \infty} \int_{\Omega_\epsilon} F(t_\epsilon w_n^+ + s_\epsilon w_n^-). \quad (3.6)$$

Now the real essence of the modification of penalization really comes up. Note that for $\epsilon > 0$ sufficiently small

$$I_{\epsilon, \mathbb{R}^N \setminus \Omega_\epsilon}(v) := \frac{1}{2} \int_{\mathbb{R}^N \setminus \Omega_\epsilon} (|\nabla v|^2 + V(\epsilon x) v^2) - \int_{\mathbb{R}^N \setminus \Omega_\epsilon} \tilde{F}_\epsilon(v)$$

is a strictly convex functional in $\tilde{H}(\mathbb{R}^N \setminus \Omega_\epsilon) = \left\{ v \in H^1(\mathbb{R}^N \setminus \Omega_\epsilon), \int_{\mathbb{R}^N \setminus \Omega_\epsilon} (|\nabla v|^2 + V(\epsilon x) v^2) < \infty \right\}$. In fact, for $v, h \in \tilde{H}(\mathbb{R}^N \setminus \Omega_\epsilon)$, $h \neq 0$, by (2.2) and Sobolev embeddings

$$\begin{aligned} I''_{\epsilon, \mathbb{R}^N \setminus \Omega_\epsilon}(v)(h, h) &= \|h\|_{\epsilon, \mathbb{R}^N \setminus \Omega_\epsilon}^2 - \int_{\mathbb{R}^N \setminus \Omega_\epsilon} \tilde{f}'_\epsilon(v) h^2 \\ &\geq \|h\|_{\epsilon, \mathbb{R}^N \setminus \Omega_\epsilon}^2 (1 - \epsilon^\tau) > 0, \end{aligned}$$

for $\epsilon > 0$ sufficiently small. Then, by convex analysis it follows that $I_{\epsilon, \mathbb{R}^N \setminus \Omega_\epsilon}$ is weakly lower semicontinuous. Then (3.5), (3.6) and this fact imply that

$$\begin{aligned} I_\epsilon(t_\epsilon w_\epsilon^+ + s_\epsilon w_\epsilon^-) &\leq \liminf_{n \rightarrow \infty} I_\epsilon(t_\epsilon w_\epsilon^+ + s_\epsilon w_\epsilon^-) \\ &\leq \liminf_{n \rightarrow \infty} (I_\epsilon(t_\epsilon w_\epsilon^+) + I_\epsilon(s_\epsilon w_\epsilon^-)) \\ &\leq \liminf_{n \rightarrow \infty} (I_\epsilon(w_\epsilon^+) + I_\epsilon(w_\epsilon^-)) \\ &= b_\epsilon. \end{aligned}$$

Hence $I_\epsilon(t_\epsilon w_\epsilon^+ + s_\epsilon w_\epsilon^-) = b_\epsilon$.

The second step is proving that $v_\epsilon \in \tilde{H}$ which minimizes I_ϵ on $\mathcal{N}_\epsilon^\pm$ is a critical point of I_ϵ in $\tilde{H}$. This can be done by employing the same arguments of Section 3 in [6]. For the sake of completeness we include all the details of this proof.

Supposing by contradiction that $I'_\epsilon(v_\epsilon) \neq 0$, there exists $\delta, \lambda > 0$ such that

$$\|I'_\epsilon(v)\|_* \geq \lambda, \quad \text{for all } v \in B_\delta(v_\epsilon) \subset \tilde{H}. \quad (3.7)$$

Let us consider the function $(t, s) \mapsto tv_\epsilon^+ + sv_\epsilon^-$ defined on $D = (\frac{1}{2}, \frac{3}{2})$ and note that by Lemma 3.2

$$\delta_\epsilon := \max_{(t,s) \in \partial D} I_\epsilon(tv_\epsilon^+ + sv_\epsilon^-) < I_\epsilon(v_\epsilon) = d_\epsilon. \quad (3.8)$$

Taking $\rho = \min \left\{ \frac{d_\epsilon - \delta_\epsilon}{2}, \frac{\lambda \delta}{24} \right\}$ and $S = B_{\delta/3}(v_\epsilon)$, Lemma 2.3 in [22] yields a deformation η such that

$$i) \quad \eta(1, v) = v \text{ for all } v \notin I_\epsilon^{-1}([d_\epsilon - 2\rho, d_\epsilon + 2\rho]),$$

$$ii) \quad \eta(1, I_\epsilon^{d_\epsilon + \rho} \cap S) \subset I_\epsilon^{d_\epsilon - \rho},$$

$$iii) \quad I_\epsilon(\eta(1, v)) \leq I_\epsilon(v), \text{ for all } v \in \tilde{H}.$$

Claim.

$$\max_{(t,s) \in \overline{D}} I_\epsilon(\eta(1, tv_\epsilon^+ + sv_\epsilon^-)) < d_\epsilon.$$

In fact, if $(t, s) \neq (1, 1)$, then by Lemma 3.2,

$$I_\epsilon(\eta(1, tv_\epsilon^+ + sv_\epsilon^-)) \leq I_\epsilon(tv_\epsilon^+ + sv_\epsilon^-) < d_\epsilon.$$

On the other hand, by *ii*),

$$I_\epsilon(\eta(1, v_\epsilon)) < d_\epsilon - \rho,$$

which proves the claim.

Now, let us prove that there exists $(t, s) \in D$ such that $\eta(1, tv_\epsilon^+ + sv_\epsilon^-) \cap \mathcal{N}_\epsilon^\pm \neq \emptyset$, which together with the Claim, contradicts the definition of d_ϵ .

Define $h(t, s) = \eta(1, tv_\epsilon^+ + sv_\epsilon^-)$ and $\Psi_0, \Psi_1 : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by

$$\Psi_0(t, s) = (I'_\epsilon(tv_\epsilon^+)v_\epsilon^+, I'_\epsilon(sv_\epsilon^-)v_\epsilon^-),$$

and

$$\Psi_1(t, s) = \left(\frac{1}{t} I'_\epsilon(h(t, s)^+)h(t, s)^+, \frac{1}{s} I'_\epsilon(h(t, s)^-)h(t, s)^- \right).$$

By results of Brower Degree Theory, $\deg(\Psi_0, D, (0, 0)) = 1$. On the other hand, note that by (3.8), $(t, s) \mapsto tv_\epsilon^+ + sv_\epsilon^-$ coincides with h on ∂D . Hence $\Psi_0 = \Psi_1$ on ∂D and then $\deg(\Psi_1, D, (0, 0)) = \deg(\Psi_0, D, (0, 0)) = 1$. Therefore there exists $(t, s) \in D$ such that $\Psi_1(t, s) = (0, 0)$ and consequently

$$\eta(1, tv_\epsilon^+ + sv_\epsilon^-) = h(t, s) \in \mathcal{N}_\epsilon^\pm.$$

Finally, this contradiction proves the theorem. $\square$

4 Concentration results

Let us introduce a sequence $\epsilon_n \rightarrow 0$ as $n \rightarrow \infty$ and, for each $n \in \mathbb{N}$, let us denote by v_n the solution v_{ϵ_n} given by Theorem 3.4 and consider $d_n := d_{\epsilon_n}$, $\|\cdot\|_n := \|\cdot\|_{\epsilon_n}$ and $I_n := I_{\epsilon_n}$.

The following result provides an upper estimate for the sequence of minimax values d_n . Its proof is inspired in the arguments of Alves and Soares in [1].

Lemma 4.1.

$$\limsup_{n \rightarrow \infty} \epsilon_n^k d_n \leq 2\omega_k \inf_{\Omega \cap \mathcal{H}^\perp} \mathcal{M}$$

and

$$\epsilon_n^k \|v_n\|_n^2 \leq C.$$

Proof. Let $z_0 = (x_0, y_0) \in \Omega$ be such that $\mathcal{M}(z_0) = \inf_{x \in \Omega} \mathcal{M}(x)$. Since Ω is an open set, there exists $R > 0$ such that $B_{2R}(z_0) \subset \Omega$. Let us choose points $z_1, z_2 \in \partial B_R(z_0)$ such that, if $z_i = (Q'_i, Q''_i)$, then $|Q''_1 - Q''_2| = 2R$.

Note that $B_R(z_i) \subset \Omega$ for $i = 1, 2$. In the rest of this proof, $i \in \{1, 2\}$. Let us choose smooth cut-off functions $\psi_i : \mathbb{R}^{N-k} \rightarrow \mathbb{R}$ such that $\psi_i = 1$ in $B_{\mathbb{R}^{N-k}}((Q'_i, |Q''_i|), R/4)$ and $\psi_i = 0$ in $\mathbb{R}^{N-k} \setminus B_{\mathbb{R}^{N-k}}((Q'_i, |Q''_i|), R/2)$.

Let $w_i \in H^1(\mathbb{R}^{N-k})$, be a ground-state solution of

$$-\Delta u + V(z_i)u = f(u) \quad \text{in } \mathbb{R}^{N-k}$$

and $w_{n,z_i} : \mathbb{R}^N \rightarrow \mathbb{R}$ be given by

$$w_{n,z_i}(x', x'') = \psi_i(\epsilon_n x', \epsilon_n |x''|) w_i \left(x' - \frac{Q'_i}{\epsilon_n}, |x''| - \frac{|Q''_i|}{\epsilon_n} \right).$$

We associate for each $w_{n,z_i}(x)$ its k -dimensional counterpart $\tilde{w}_{n,z_i}(x', r) = w_{n,z_i}(x', x'')$ where $|x''| = r \in \mathbb{R}$. It is well known the existence of $t_{i_n} > 0$ such that $t_{i_n} w_{n,z_i} \in \mathcal{N}_{\epsilon_n}$. By the construction we have made so far, it is straightforward to see that

$$\overline{w}_n := t_{1_n} w_{n,z_1} - t_{2_n} w_{n,z_2} \in \mathcal{N}_{\epsilon_n}^{\pm}.$$

Using the fact that I_n is even, we have

$$d_n \leq I_n(\overline{w}_n) = I_n(t_{1_n} w_{n,z_1}) + I_n(t_{2_n} w_{n,z_2}). \quad (4.1)$$

By a change of variable, let us note that for $i \in \{1, 2\}$, $I_n(t_n w_{n,z_i})$ is equal

to

$$\begin{aligned}
& \frac{t_{i_n}^2}{2} \int_{\mathbb{R}^N} (|\nabla w_{n,z_i}|^2 + V(\epsilon_n x) w_{n,z_i}^2) dx - \int_{\mathbb{R}^N} G_n(\epsilon_n x, t_{i_n} w_{n,z_i}) dx \\
&= \omega_k \left[\frac{t_{i_n}^2}{2} \int_{\mathbb{R}^{N-k-1}} \int_0^{+\infty} r^k (|\nabla \tilde{w}_{n,z_i}(x', r)|^2 + V(\epsilon_n x', \epsilon_n r) \tilde{w}_{n,z_i}(x', r)^2) dr dx' \right. \\
&\quad \left. - \int_{\mathbb{R}^{N-k-1}} \int_0^{+\infty} r^k G_n(\epsilon_n x, \epsilon_n r, t_{i_n} \tilde{w}_{n,z_i}(x', r)) dr dx' \right] \\
&= \omega_k \left[\frac{t_{i_n}^2}{2} \int_{\mathbb{R}^{N-k-1}} \int_{\frac{-|Q_i''|}{\epsilon_n}}^{+\infty} \left(\sigma + \frac{|Q_i''|}{\epsilon_n} \right)^k (|\nabla(\psi_i(\epsilon_n x' + Q_i', \epsilon_n \sigma + |Q_i''|) w_i(x', \sigma))|^2 \right. \\
&\quad \left. + V(\epsilon_n x' + \epsilon_n Q_i', \epsilon_n \sigma + |Q_i''|) (\psi_i(\epsilon_n x' + \epsilon_n Q_i', \epsilon_n \sigma + |Q_i''|) w_i(x', \sigma))^2) d\sigma dx' \right. \\
&\quad \left. - \int_{\frac{-|Q_i''|}{\epsilon_n}}^{+\infty} \left(\sigma + \frac{|Q_i''|}{\epsilon_n} \right)^k G_n(\epsilon_n x' + \epsilon_n Q_i', \epsilon_n \sigma + |Q_i''|, t_{i_n} \psi_i(\epsilon_n x' + \epsilon_n Q_i', \epsilon_n \sigma + |Q_i''|) w_i(x', \sigma)) d\sigma dx' \right] \\
&= \epsilon_n^{-k} \omega_k |Q_i|^k I_{V(z_i)}(t_{i_n} w_i) + o(1) \leq \epsilon_n^{-k} \omega_k |Q_i|^k \mathcal{E}(V(z_i)) = \epsilon_n^{-k} \omega_k \mathcal{M}(z_i) + o(1).
\end{aligned}$$

By the last inequality and (4.1) we have

$$\epsilon_n^k d_n \leq \omega_k (\mathcal{M}(z_1) + \mathcal{M}(z_2)) + o(1) \quad (4.2)$$

and making $R \rightarrow 0$, continuity of V and $\mathcal{M}$ implies in the result.

Now, in order to see that

$$\epsilon_n^k \|v_n\|_n^2 \leq C,$$

just observe that $v_n \in \mathcal{N}_{\epsilon_n}$ and use *ii*) of Lemma 3.5. $\square$

The next lemma implies that solutions found in Theorem 3.4 do not vanish when $n \rightarrow \infty$.

Lemma 4.2. *Let P_n^1, P_n^2 be local maximum and minimum points of v_n , respectively. Then $P_n^i \in \Omega_{\epsilon_n} := \epsilon_n^{-1} \Omega$,*

$$v_n(P_n^1) \geq a \quad \text{and} \quad v_n(P_n^2) \leq -a,$$

where $a > 0$ is such that $f(a)/a = V_0/2$.

Proof. First of all let us prove that $P_n^i \in \Omega_{\epsilon_n}$. Suppose by contradiction that $P_n^1 \notin \Omega_{\epsilon_n}$. Since P_n^1 is a local maximum point, it follows that $\Delta v_n(P_n^1) \leq 0$. By definition of g_n we have

$$V_0 v_n(P_n^1) \leq -\Delta v_n(P_n^1) + V(\epsilon_n P_n^1) v_n(P_n^1) = \tilde{f}_{\epsilon_n}(v_n(P_n^1)) \leq \epsilon_n^\tau v_n(P_n^1),$$

which is impossible for $\epsilon > 0$ sufficiently small. A similar argument applies to P_n^2 .

Since $P_n^1 \in \Omega_{\epsilon_n}$ and $v_n(P_n^1) > 0$, from the definition of g_n we have

$$0 \geq \Delta v_n(P_n^1) = \left(V(\epsilon_n P_n^1) - \frac{f(v_n(P_n^1))}{v_n(P_n^1)} \right) v_n(P_n^1).$$

Supposing by contradiction that $v_n(P_n^1) < a$. By the choice of $a > 0$ and (f_5) it follows that

$$V_0 \leq V(\epsilon_n P_n^1) \leq \frac{f(v_n(P_n^1))}{v_n(P_n^1)} \leq \frac{f(a)}{a} = \frac{V_0}{2}, \quad (4.3)$$

which is a contradiction. Analogously we prove that $v_n(P_n^2) \leq -a$. $\square$

By the last result, there exist $P^1, P^2 \in \Omega$ such that, along a subsequence

$$\lim_{n \rightarrow \infty} \epsilon_n P_n^i = P^i, \quad i \in \{1, 2\}. \quad (4.4)$$

The same argument of [1] with short modifications can be used to prove the following result.

Lemma 4.3. *Using the same notation that in the last result, it follows that*

$$\lim_{n \rightarrow \infty} |P_n^1 - P_n^2| = +\infty. \quad (4.5)$$

Lemma 4.4. *Let $y_n = (y'_n, y''_n) \subset \mathbb{R}^N$ be a sequence such that $\epsilon_n y_n \rightarrow (\bar{y}', \bar{y}'') \in \bar{\Omega}$ as $n \rightarrow \infty$. Denoting $\tilde{v}_n(x', r) := v_n(x', x'')$ where $|x''| = r$, let us define $\tilde{w}_n : \mathbb{R}^{N-k-1} \times [-|y_n|, +\infty) \rightarrow \mathbb{R}$ by*

$$\tilde{w}_n(x', r) := \tilde{v}_n(x', r + |y''_n|).$$

Then there exists $\tilde{w} \in H^1(\mathbb{R}^{N-k})$ such that $\tilde{w}_n \rightarrow \tilde{w}$ in $C_{loc}^2(\mathbb{R}^{N-k})$ and $\tilde{w}$ satisfies the limit problem

$$-\Delta \tilde{w} + V(\bar{y}', |\bar{y}''|) \tilde{w} = \tilde{g}_n(x', r, \tilde{w}) \quad \text{in } \mathbb{R}^{N-k}, \quad (4.6)$$

where $\tilde{g}_n(x', r, s) := \chi(x', r) f(s) + (1 - \chi(x', r)) \tilde{f}_{\epsilon_n}(s)$ and $\chi(x', r) = \lim_{n \rightarrow \infty} \chi_\Omega(\epsilon_n x' + \epsilon_n y'_n, \epsilon_n r + \epsilon_n |y''_n|)$.

Proof. The proof is analogous to [7][Lemma 4.3] but we sketch it here for the sake of completeness.

Note that $\tilde{w}_n$ satisfies the following problem

$$-\Delta \tilde{w}_n - \frac{k}{(r + |y_n''|)} \frac{\tilde{w}_n}{\partial r} + V(\epsilon_n x' + \epsilon_n y_n', \epsilon_n r + \epsilon_n |y_n''|) \tilde{w}_n = g_n(\epsilon_n x' + \epsilon_n y_n', \epsilon_n r + \epsilon_n |y_n''|, \tilde{w}_n), \quad (4.7)$$

in $\mathbb{R}^{N-k-1} \times [-|y_n|, +\infty)$.

By Lemma 4.1 it follows that

$$\int_{\mathbb{R}^{N-k-1}} \int_{-|y_n''|}^{+\infty} (|\nabla \tilde{w}_n|^2 + V(\epsilon_n x' + \epsilon_n y_n', \epsilon_n r + \epsilon_n |y_n''|) \tilde{w}_n^2) dr dx' \leq C, \quad (4.8)$$

uniformly in n and then, for some $\tilde{w} \in H^1(\mathbb{R}^{N-k})$,

$$\tilde{w}_n \rightharpoonup \tilde{w} \quad \text{in } H^1(\mathbb{R}^{N-k}). \quad (4.9)$$

By choosing a sequence $R_n \rightarrow \infty$ such that $\epsilon_n R_n \rightarrow 0$ and considering a smooth cut-off function in $\mathbb{R}^{N-k}$, η_R such that $0 \leq \eta_R \leq 1$, $\eta_R(z) = 1$ if $|z| \leq \frac{R}{2}$ and $\eta_R(z) = 0$ if $|z| > R$, and $\|\nabla \eta_R\|_\infty \leq \frac{C}{R}$, it can be proved using (4.8) that $\bar{w}_n(z) := \eta_{R_n}(z) \tilde{w}_n(z)$ is bounded in $H^1(\mathbb{R}^{N-k})$, uniformly in n .

Since $\bar{w}_n$ satisfies (4.7) in $B(0, R_n)$, it follows by classical elliptic estimates that

$$\|\bar{w}_n\|_{W^{2,q}(B(0,R))} \leq C \quad (4.10)$$

for sufficiently large $n \in \mathbb{N}$, where $R > 0$ is fixed.

By (4.9) and (4.10) it follows that $\tilde{w}_n \rightarrow \tilde{w}$ in $C_{loc}^2(\mathbb{R}^{N-k})$ and that $\tilde{w}$ satisfies (4.6). $\square$

Since the concentration set is expected to be a sphere in $\mathbb{R}^N$, it is natural to introduce the distance between two k -dimensional spheres in $\mathbb{R}^N$, which gives rise to neighborhoods in which we want to estimate the mass of solutions. For $x = (x', x''), y = (y', y'') \in \mathbb{R}^N$, let

$$d_k(x, y) = \sqrt{(x' - y')^2 + (|x''| - |y''|)^2}, \quad (4.11)$$

which corresponds to the distance between k -dimensional spheres centered at the origin, parallel to $\mathcal{H}^\perp$ and of radius $|x''|$ and $|y''|$, respectively. According to this distance, the balls are given by

$$B_k(x, r) = \{y \in \mathbb{R}^N; d_k(x, y) < r\}.$$

From now on, for $\Lambda \subset \mathbb{R}^N$ we denote

$$I_{n,\Lambda}(v) = \frac{1}{2} \int_{\Lambda} (|\nabla v|^2 + V(\epsilon_n x) v^2) dx - \int_{\Lambda} G_n(\epsilon_n x, v) dx.$$

The following is the main step in proving the concentration result.

Proposition 4.5. *Suppose all the assumptions of Theorem 1.1 to hold. Then,*

$$i) \quad \lim_{n \rightarrow \infty} \epsilon_n^k d_n = 2\omega_k \inf_{\Omega} \mathcal{M}.$$

$$ii) \quad \lim_{n \rightarrow \infty} \mathcal{M}(\epsilon_n P_n^i) = \inf_{\Omega} \mathcal{M}, \quad i \in \{1, 2\}.$$

Proof. Let us get started with $i)$. By Lemma 4.1 it follows that

$$\limsup_{n \rightarrow \infty} \epsilon_n^k d_n \leq 2\omega_k \inf_{\Omega} \mathcal{M}. \quad (4.12)$$

In order to prove that

$$\liminf_{n \rightarrow \infty} \epsilon_n^k d_n \geq 2\omega_k \inf_{\Omega} \mathcal{M}, \quad (4.13)$$

we use some ideas of [12][Lemma 2.2].

By Lemmas 4.2 and 4.4 it follows that $\tilde{w}_n^i(x', r) := \tilde{v}_n(x', r + P_n^i) \rightarrow \tilde{w}_i$ in $C_{loc}^2(\mathbb{R}^{N-k})$, where $\tilde{w}_i \neq 0$ and satisfies (4.6) with $(\tilde{y}', \tilde{y}'') = P^i$.

For each $R > 0$ and up to a subsequence in n , Lemma 4.4 with calculations similar to which have resulted in (4.2) implies that

$$\begin{aligned} \epsilon_n^k I_{n, B_k(P^i, R)}(v_n) &= \omega_k |P^{i''}|^k \left(\frac{1}{2} \int_{B_R(0)} (|\nabla \tilde{w}_i|^2 + V(P^i) \tilde{w}_i^2) dz - \right. \\ &\quad \left. \int_{B_R(0)} \tilde{G}(r, \tilde{w}_i) dz \right) + o_n(1) \\ &\geq \omega_k |P^{i''}|^k \left(\frac{1}{2} \int_{B_R(0)} (|\nabla \tilde{w}_i|^2 + V(P^i) \tilde{w}_i^2) dz - \right. \\ &\quad \left. \int_{B_R(0)} F(\tilde{w}_i) dz \right) + o_n(1). \end{aligned}$$

Since $\tilde{w}_i \in H^1(\mathbb{R}^{N-k})$, it follows that for a given $\eta > 0$, there exists $R > 0$ such that

$$\liminf_{n \rightarrow \infty} \epsilon_n^k I_{n, B_k(y_n, R)}(v_n) \geq \omega_k \mathcal{M}(P^i) - \eta.$$

Taking into account Lemma 4.3, it follows that for n large enough $B_k(P_n^1, R)$ and $B_k(P_n^2, R)$ are disjoint and then

$$\epsilon_n^k I_n(v_n) = \epsilon_n^k I_{n, B_k(P_n^1, R)}(v_n) + \epsilon_n^k I_{n, B_k(P_n^2, R)}(v_n) + I_{n, \mathbb{R}^N \setminus B_{n, R}}(v_n),$$

where $B_{n, R} := B_k(P_n^1, R) \cup B_k(P_n^2, R)$.

Now, calculations similar to (2.21) in [17] implies that $\liminf_{n \rightarrow \infty} I_{n, \mathbb{R}^N \setminus B_{n, R}}(v_n) \geq -\eta$ for $R > 0$ sufficiently large and then

$$\begin{aligned} \liminf_{n \rightarrow \infty} \epsilon_n^k I_n(v_n) &\geq \omega_k (\mathcal{M}(P^1) + \mathcal{M}(P^2)) - 3\eta \\ &\geq 2\omega_k \inf_{\Omega} \mathcal{M} - 3\eta. \end{aligned}$$

Since the last inequality holds for all $\eta > 0$, (4.13) is proved.

To prove *ii*) let us suppose by contradiction that

$$\mathcal{M}(P^i) > \inf_{\Omega} \mathcal{M}, \quad i \in \{1, 2\}.$$

where P^i is given by (4.4). Just by arguing as in the first item, one can see that

$$\liminf_{n \rightarrow \infty} \epsilon_n^k I_n(v_n) \geq \omega_k (\mathcal{M}(P^1) + \mathcal{M}(P^2)) > 2\omega_k \inf_{\Omega} \mathcal{M},$$

which contradicts the statement of Lemma 4.1. This contradiction proves the proposition. $\square$

5 Proof of Theorem 1.1

Let v_n be as in the beginning of Section 4 and $u_n(x) := v_n(\epsilon_n^{-1}x)$.

Proposition 5.1.

$$\lim_{n \rightarrow \infty} \|v_n\|_{L^\infty(\Omega_n \setminus (B_k(P_n^1, R) \cup B_k(P_n^2, R)))} = 0.$$

Proof. The prove follows by contradiction. Let us suppose that there exist $\eta > 0$ and sequences $R_n \rightarrow \infty$ and $(y_n) \subset \Omega_n \setminus (B_k(P_n^1, R_n) \cup B_k(P_n^1, R_n))$ such that

$$|v_n(y_n)| \geq \eta.$$

Since $\epsilon_n y_n \in \Omega$, it follows that $\epsilon_n y_n \rightarrow \bar{y} \in \Omega$ up to a subsequence.

Then following the arguments in Proposition 4.5 it is possible to show that

$$\epsilon_n^k d_n = I_n(v_n) \geq 3\omega_k \inf_{\Omega} \mathcal{M},$$

which contradicts Lemma 4.1. $\square$

By standard elliptic regularity theory it is possible to show that $v_n \in C^2(\mathbb{R}^N)$. Then, by continuity, Proposition 5.1 implies that

$$\|v_n\|_{L^\infty(\partial((B_k(P_n^1, R_n) \cup B_k(P_n^1, R_n)))} = o_n(1) \quad (5.1)$$

In order to prove the exponential decay with respect to ϵ_n of the functions u_ϵ , let us take as a comparison function

$$W(x) = C(e^{-\beta d_k(x, P_n^1)} + e^{-\beta d_k(x, P_n^2)}),$$

defined in $\mathbb{R}^N \setminus (B_k(P_n^1, R) \cup B_k(P_n^1, R))$, where $C > 0$ is to be choosen. For sufficiently small $\beta > 0$ (independent of ϵ_n), it follows that

$$\left(-\Delta + V(\epsilon_n x) - \frac{g_n(\epsilon_n x, v_n)}{v_n} \right) (W \pm v_n) \geq 0 \quad \text{in } \mathbb{R}^N \setminus (B_k(P_n^1, R) \cup B_k(P_n^1, R)).$$

Then, by (5.1), for $x \in \partial(B_k(P_n^1, R) \cup B_k(P_n^1, R))$

$$W(x) \pm v_n(x) = C2e^{-\beta R} \pm v_n \geq 0$$

for a sufficiently large constant $C > 0$ which does not depend on n . Hence Maximum Principle applies and

$$|v_n| \leq W(x), \quad \text{on } \mathbb{R}^N \setminus (B_k(P_n^1, R) \cup B_k(P_n^1, R)).$$

Then there exists a sufficiently large constant $C > 0$ such that

$$|v_n(x)| \leq C \left(e^{-\beta d_k(x, P_n^1)} + e^{-\beta d_k(x, P_n^2)} \right),$$

for $x \in \mathbb{R}^N \setminus (B_k(P_n^1, R) \cup B_k(P_n^1, R))$, which implies that

$$|u_n(x)| \leq C \left(e^{-\frac{\beta}{\epsilon_n} d_k(x, \epsilon_n P_n^1)} + e^{-\frac{\beta}{\epsilon_n} d_k(x, \epsilon_n P_n^2)} \right) \quad (5.2)$$

for $x \in \mathbb{R}^N \setminus (B_k(\epsilon_n P_n^1, \epsilon_n R) \cup B_k(\epsilon_n P_n^1, \epsilon_n R))$. In particular it holds that

$$\|u_n\|_{L^\infty(\mathbb{R}^N \setminus \Omega)} \leq C e^{\frac{-\beta}{\epsilon_n}}$$

which implies that u_n satisfies the original problem. The proof of (1.5) follows by (4.4) and item *ii*) of Proposition 4.5. Therefore the proof of Theorem 1.1 follows.

Acknowledgment.

We would like to express our gratitude to Prof. Antonio Suárez for some discussions on this subject.

Competing interests. The authors declare that they have no competing interests.

Author's contribution. All the author wrote, review and approve the final manuscript.

References

- [1] C.O. Alves, S.H.M. Soares *On the location and profile of spike-layer nodal solutions to nonlinear Schrödinger equations*. Journal of Mathematical Analysis and Applications **296** (2004), 563 - 577.
- [2] C.O. Alves, S.H.M. Soares, *Nodal solutions for singularly perturbed equations with critical exponential growth*, Journal of Differential Equations **234** (2007), 464 - 484.
- [3] C.O. Alves, M. Souto, *Existence of least energy nodal solution for a Schrödinger-Poisson system in bounded domains*, to appear in Z. Angew. Math. Phys.
- [4] A. Ambrosetti, A. Malchiodi, W. Ni *Singularly perturbed elliptic equations with symmetry: Existence of solutions concentrating on spheres, Part I*, Commun. Math. Phys., **235** (2003), 427 - 466.

- [5] A. Ambrosetti, A. Malchiodi, W. Ni *Singularly perturbed elliptic equations with symmetry: Existence of solutions concentrating on spheres, Part II*, Indiana Univ. Math. J., **53**, No. 2, (2004), 297 - 329.
- [6] T. Bartsch, T. Weth, M. Willem, *Partial symmetry of least energy nodal solutions to some variational problems*, J. Anal. Math. **96** (2005), 1 - 18.
- [7] D. Bonheure, J. Di Cosmo, J.V. Schaftingen, *Nonlinear Schrödinger equation with unbounded or vanishing potentials: solutions concentrating on lower dimensional spheres*, Journal of Differential Equations. **252** (2012), 941 - 968.
- [8] D. Bonheure, J. Di Cosmo, C. Mercuri, *Concentration on circles for nonlinear Schrödinger-Poisson systems with unbounded potentials vanishing at infinity*, Commun. Contemp. Math. **14**, No. 2 (2012), 1 - 30.
- [9] A. Castro, J. Cossio, J. Neuberger *A sign-changing solution for a super linear Dirichlet problem*, Rocky Mountain J. of Math. **27** (4) (1997), 1041-1053.
- [10] S. Cingolani, A. Pistoia, *Nonexistence of single blow-up solutions for a nonlinear Schrödinger equation involving critical Sobolev exponent*, Z. Angew. Math. Phys. **55** (2004), 201 - 215.
- [11] T. D'Aprile, W. Juncheng, *On bound states concentrating on spheres for the Maxwell-Schrödinger equation*, SIAM J. Math. Anal. **37** (2005), 321 - 342.
- [12] M. del Pino, P. Felmer, *Local mountain pass for semilinear elliptic problems in unbounded domains*, Calc. Var. Partial Differential Equations **4** (1996), 121 - 137.
- [13] A. Floer, A. Weinstein, *Nonspreading wave packets for the cubic Schrödinger equations with a bounded potential*, Journal of Functional Analysis **69** (1986), 397 - 408.
- [14] M. Fei, *Sign-changing multi-peak solutions for nonlinear Schrödinger equations with compactly supported potential*, Acta Appl. Math. **127** (2013), 137 - 154.

- [15] I. Ianni, *Solutions of the Schrödinger-Poisson problem concentrating on spheres, Part II: existence*, Math. Models Methods Appl. Sci. **19**, No. 6 (2009), 877 - 910.
- [16] A. Malchiodi, M. Montenegro, *Boundary concentration phenomena for a singularly perturbed elliptic problem*, Commun. Pure Appl. Math **55** (2002), 1507 - 1568.
- [17] M. Pimenta, S. Soares, *Existence and concentration of solutions for a class of biharmonic equations*, J. Math. Anal. Appl. **390** (2012), 274 - 289.
- [18] P.H. Rabinowitz, *On a class of nonlinear Schrödinger equations*, ZAMP **43** (1992), 270 - 291.
- [19] D. Ruiz, *Semiclassical states for coupled Schrödinger-Maxwell equations: concentration around a sphere*, Math. Models Methods Appl. Sci. **15** (2005), 141 - 164.
- [20] Y. Sato, *Sign-changing multi-peak solutions for nonlinear Schrödinger equations with critical frequency*, Comm. Pure Appl. Anal. **7**, N.4, (2008), 883 - 903.
- [21] X. Wang, *On concentration of positive bound states of nonlinear Schrödinger equations*, Commun. Math. Phys. **153** (1993), 229 - 244.
- [22] M. Willem, *Minimax theorems*, Birkhäuser Boston (1996).